

Immunization Requirements

DISEASE(S)	MEETS IMMUNIZATION REQUIREMENTS	COMMENTS
DTaP	(AGE 1-6 YEARS): 4 doses, with one dose given on or after the 4th birthday, OR any 5 doses. (AGE 7-9 YEARS): 3 doses of Td or any previously administered combination of DTP, DTaP, and DT to equal 3 doses.	Any child entering pre-school, pre-Kindergarten, or Kindergarten needs a minimum of four doses. Pupils after the seventh birthday should receive adult type Td. DTP/Hib vaccine and DTaP also valid DTP doses. Laboratory evidence of immunity is also acceptable.
Tdap	GRADE 6 (<i>or comparable age level for special education programs</i>): 1 dose	For pupils entering Grade 6 on or after 9-1-08 and born on or after 1-1-97. A child does not need a Tdap dose until FIVE years after the last DTP/DTaP or Td dose.
POLIO	(AGE 1-6 YEARS): 3 doses, with one dose given on or after the 4th birthday, OR any 4 doses. (AGE 7 or OLDER): Any 3 doses.	Either Inactivated Polio Vaccine (IPV) or Oral Polio Vaccine (OPV) separately or in combination is acceptable. Polio vaccine is not required of pupils 18 years of age or older. Laboratory evidence of immunity is also acceptable.
MEASLES	If born before 1-1-90, 1 dose of a live Measles-containing vaccine. If born on or after 1-1-90, 2 doses of a live Measles-containing vaccine. If entering a college or university after 9-1-95 and previously unvaccinated, 2 doses of a live Measles-containing vaccine.	Any child over 15 months of age entering child care, pre-school, or pre-Kindergarten needs a minimum of 1 dose of measles vaccine. Any child entering Kindergarten needs 2 doses. Previously unvaccinated students entering college after 9-1-95 need 2 doses of measles-containing vaccine or any combination containing live measles virus administered after 1968. Documentation of 2 prior doses is acceptable. Laboratory evidence of immunity is also acceptable. Intervals between first and second measles/MMR/MR doses cannot be less than 1 month.
RUBELLA and MUMPS	1 dose of live Mumps-containing vaccine. 1 dose of live Rubella-containing vaccine.	Any child over 15 months of age entering child care, pre-school, or pre-Kindergarten needs 1 dose of rubella and mumps vaccine. Each student entering college for the first time after 9-1-95 needs 1 dose of rubella and mumps vaccine or any combination containing live rubella and mumps virus administered after 1968. Laboratory evidence of immunity is also acceptable.
VARICELLA	1 dose on or after first birthday.	All children 19 months of age and older enrolled into a child care/pre-school center after 9-1-04 or children born on or after 1-1-98 entering a school for the first time in Kindergarten or Grade 1 need 1 dose of varicella vaccine. Laboratory evidence of immunity, physician's statement or a parental statement of previous varicella disease is also acceptable.

Immunization Requirements

<p>HAEMOPHILUS INFLUENZAE B (Hib)</p>	<p>(AGE 2-11 MONTHS)⁽¹⁾: 2 doses (AGE 12-59 MONTHS)⁽²⁾: 1 dose</p>	<p>Mandated only for children enrolled in child care, pre-school, or pre-Kindergarten. ⁽¹⁾ Minimum of 2 doses of Hib vaccine is needed if between the ages of 2-11 months. ⁽²⁾ Minimum of 1 dose of Hib vaccine is needed after the first birthday. DTP/Hib and Hib/Hep B also valid Hib doses.</p>
<p>HEPATITIS B</p>	<p>(Pre-K-GRADE 12): 3 doses or 2 doses ⁽¹⁾</p>	<p>⁽¹⁾ If a child is between 11-15 years of age and has not received 3 prior doses of Hepatitis B then the child is eligible to receive 2-dose Hepatitis B Adolescent formulation. Laboratory evidence of immunity is also acceptable.</p>
<p>PNEUMOCOCCAL</p>	<p>(AGE 2-11 MONTHS)⁽¹⁾: 2 doses (AGE 12-59 MONTHS)⁽²⁾: 1 dose</p>	<p>Mandated only for children enrolled in child care, pre-school, or pre-Kindergarten. ⁽¹⁾ Minimum of 2 doses of Pneumococcal vaccine is needed if between the ages of 2-11 months. ⁽²⁾ Minimum of 1 dose of Pneumococcal vaccine is needed after the first birthday.</p>
<p>MENINGOCOCCAL</p>	<p>(Entering GRADE 6 <i>(or comparable age level for Special Ed programs)</i>): 1 dose ⁽¹⁾ (Entering a four-year college or University, previously unvaccinated and residing in a campus dormitory): 1 dose ⁽²⁾</p>	<p>⁽¹⁾ For pupils entering Grade 6 on or after 9-1-08 and born on or after 1-1-97. ⁽²⁾ Previously unvaccinated students entering a four-year college or university after 9-1-04 and who reside in a campus dormitory, need 1 dose of meningococcal vaccine. Documentation of one prior dose is acceptable.</p>
<p>INFLUENZA</p>	<p>(AGES 6-59 MONTHS): 1 dose ANNUALLY</p>	<p>For children enrolled in child care, pre-school or pre-Kindergarten on or after 9-1-08. 1 dose to be given between September 1 and December 31 of each year.</p>